

NAME: _____

PUID: _____

ECE 656 Exam 1: Fall 2017

September 21, 2017

Mark Lundstrom
Purdue University

This is a closed book exam. You may use a calculator and the formula sheet at the end of this exam. **Only the Texas Instruments TI-30X IIS scientific calculator is allowed.**

There are four equally weighted questions. To receive full credit, you must **show your work** (scratch paper is attached).

The exam is designed to be taken in 75 minutes.

Be sure to fill in your name and Purdue student ID at the top of the page.

DO NOT open the exam until told to do so, and stop working immediately when time is called.

The last page is an equation sheet, which you may remove, if you want.

100 points possible, 25 per question

- | | | | |
|---------------|--------------|---------------|---------------|
| 1a) 20 points | 2) 25 points | 3a) 5 points | 4a) 5 points |
| 1b) 5 points | | 3b) 10 points | 4b) 10 points |
| | | 3c) 10 points | 4c) 10 points |

----- Course policy -----

I understand that if I am caught cheating in this exam, I will earn an F for the course and be reported to the Dean of Students.

Read and understood: _____
signature

- 1) In the next part of the course, we will encounter a quantity:

$$M_{3D}(E) = \frac{m^*}{2\pi\hbar^2} (E - E_c) H(E - E_c),$$

where $H(E - E_c) = 0$ for $E < E_c$ and $H(E - E_c) = 1$ for $E \geq E_c$

This problem concerns the integral: $I_1 = \int_{E_c}^{\infty} M_{3D}(E) f_0(E) dE$, where $f_0(E)$ is the equilibrium Fermi function.

- 1a) Work out the integral in terms of Fermi-Dirac integrals. **Draw a box around your answer.**

- 1b) Simplify your answer for non-degenerate (Maxwellian) statistics. **Draw a box around your answer.**

- 2) This problem concerns an integral slightly different from that in problem 1):

$$I_2 = \int_{E_c}^{\infty} M_{3D}(E) (-\partial f_0 / \partial E) dE .$$

Work out the integral in terms of Fermi-Dirac integrals. **Draw a box around your answer.**

3) The definition of the density-of-states for a 2D semiconductor is

$$D_{2D}(E) \equiv \frac{1}{A} \sum_{\vec{k}_{\parallel}} \delta(E - E_{k_{\parallel}})$$

where \vec{k}_{\parallel} is a wavevector in the plane with a magnitude of $k_{\parallel} = \sqrt{k_x^2 + k_y^2}$. The conduction band of graphene is described by


$$E_{k_{\parallel}} = \hbar v_F k_{\parallel} \quad E_{k_{\parallel}} > 0$$

3a) First, convert the sum over \vec{k}_{\parallel} to an integral in k-space.

3b) Use the graphene dispersion, $E_{k_{\parallel}} = \hbar v_F k_{\parallel}$, to convert to an integral in energy space.

3c) Evaluate the integral to find $D_{2D}(E)$ for the conduction band of graphene. Assume now that the valley degeneracy is 2 for graphene.

- 4) The material, $\text{In}_{0.53}\text{Ga}_{0.47}\text{As}$ is an important semiconductor because it is lattice matched to InP. It is a direct bandgap material with a bandgap of $E_{\text{GT}} = 0.75$ eV and an effective mass of $m_n^*/m_0 = 0.041$. It contains heavy mass, upper valleys located at an energy of $\Delta E_{\Gamma-L} = 0.55$ eV above the Γ valley minimum. You may assume that $\text{In}_{0.53}\text{Ga}_{0.47}\text{As}$ has a density-of-state effective mass in the upper valleys of $m_{\text{DOS}}^*/m_0 = 1$ and that $\text{In}_{0.53}\text{Ga}_{0.47}\text{As}$ has an optical phonon energy of 32 meV. The following two questions concern electron scattering in **undoped** $\text{In}_{0.53}\text{Ga}_{0.47}\text{As}$ at room temperature.
- 4a) Identify the important phonon scattering mechanisms in undoped $\text{In}_{0.53}\text{Ga}_{0.47}\text{As}$ at room temperature
- 4b) Sketch the total scattering rate vs. energy for **electrons in the Γ valley** from $E = 0$ (bottom of the Γ valley) to $E = 0.9$ eV. Your sketch should identify **each** of the main phonon scattering processes and the critical energies.



- 4c) Consider an electron in an L-valley with an energy $E = 0.6$ eV that scatters by optical phonon emission to a state in either the Γ -valley or one of the other L-valleys. Find the ratio of the two scattering rates:

$$\frac{1/\tau_{L \rightarrow \Gamma}}{1/\tau_{L \rightarrow L}}$$

You may assume that the deformation potentials for the two inter-valley processes are identical. **A numerical answer is required.** Note that $E = 0$ is the bottom of the Γ -valley.

SCRATCH PAPER

ECE-656 Key Equations (Weeks 1-4)

Physical constants:

$$\begin{aligned} \hbar &= 1.055 \times 10^{-34} \text{ [J-s]} & m_0 &= 9.109 \times 10^{-31} \text{ [kg]} \\ k_B &= 1.380 \times 10^{-23} \text{ [J/K]} & q &= 1.602 \times 10^{-19} \text{ [C]} & \epsilon_0 &= 8.854 \times 10^{-14} \text{ [F/cm]} \end{aligned}$$

Density of states in k-space:

$$\begin{aligned} 1D: N_k &= 2 \times (L/2\pi) = L/\pi & 2D: N_k &= 2 \times (A/4\pi^2) = A/2\pi^2 & 3D: \\ N_k &= 2 \times (\Omega/8\pi^2) = \Omega/4\pi^3 \end{aligned}$$

Density of states in energy (parabolic bands, per length, area, or volume):

$$D_{1D}(E) = \frac{g_v}{\pi \hbar} \sqrt{\frac{2m^*}{E - \epsilon_1}} \quad D_{2D}(E) = g_v \frac{m^*}{\pi \hbar^2} \quad D_{3D}(E) = g_v \frac{m^* \sqrt{2m^*(E - E_C)}}{\pi^2 \hbar^3}$$

Fermi function and Fermi-Dirac Integrals:

$$\begin{aligned} f_0(E) &= \frac{1}{1 + e^{(E - E_F)/k_B T}} \\ \mathcal{F}_j(\eta_F) &= \frac{1}{\Gamma(j+1)} \int_0^\infty \frac{\eta^j d\eta}{1 + e^{\eta - \eta_F}} & \mathcal{F}_j(\eta_F) &\rightarrow e^\eta \quad \eta_F \ll 0 & \frac{d\mathcal{F}_j}{d\eta_F} &= \mathcal{F}_{j-1} \\ \Gamma(n) &= (n-1)! \quad (n \text{ an integer}) & \Gamma(1/2) &= \sqrt{\pi} & \Gamma(p+1) &= p\Gamma(p) \end{aligned}$$

Scattering:

$$\begin{aligned} S(\vec{p}, \vec{p}') &= \frac{2\pi}{\hbar} |H_{\vec{p}', \vec{p}}|^2 \delta(E' - E - \Delta E) & H_{\vec{p}', \vec{p}} &= \frac{1}{\Omega} \int_{-\infty}^{+\infty} e^{-i\vec{p}' \cdot \vec{r}/\hbar} U_S(\vec{r}) e^{i\vec{p} \cdot \vec{r}/\hbar} d\vec{r} \\ \frac{1}{\tau(\vec{p})} &= \sum_{\vec{p}', \uparrow} S(\vec{p}, \vec{p}') & \frac{1}{\tau_m(\vec{p})} &= \sum_{\vec{p}', \uparrow} S(\vec{p}, \vec{p}') \frac{\Delta p_z}{p_{z0}} & \frac{1}{\tau_E(\vec{p})} &= \sum_{\vec{p}', \uparrow} S(\vec{p}, \vec{p}') \frac{\Delta E}{E_0} \\ \text{ADP: } |K_q|^2 &= q^2 D_A^2 & \text{ODP: } |K_q|^2 &= D_0^2 & \text{PZ: } |K_q|^2 &= (e e_{PZ} / \kappa_S \epsilon_0)^2 & \text{POP: } |K_\beta|^2 &= \frac{\rho e^2 \omega_0^2}{q^2 \kappa_0 \epsilon_0} \left(\frac{\kappa_0}{\kappa_\infty} - 1 \right) \\ S(\vec{p}, \vec{p}') &= \frac{\pi}{\Omega \rho \omega} |K_q|^2 \left(N_\omega + \frac{1}{2} \mp \frac{1}{2} \right) \delta(\vec{p}' - \vec{p} \mp \hbar \vec{q}) \delta(E' - E \mp \hbar \omega) \\ \delta(\vec{p}' - \vec{p} \mp \hbar \vec{q}) \delta(E' - E \mp \hbar \omega) &\rightarrow \frac{1}{\hbar v q} \delta \left(\pm \cos \theta + \frac{\hbar q}{2p} \mp \frac{\omega_q}{v q} \right) \\ \frac{1}{\tau} &= \frac{1}{\tau_{abs}} + \frac{1}{\tau_{ems}} = \frac{2\pi}{\hbar} \left(\frac{D_A^2 k_B T}{c_i} \right) \frac{D_{3D}(E)}{2} \quad (\text{ADP}) & L_D &= \sqrt{\frac{\kappa_S \epsilon_0 k_B T}{q^2 n_0}} \\ \frac{1}{\tau} &= \frac{1}{\tau_m} = \frac{2\pi}{\hbar} \left(\frac{\hbar D_0^2}{2\rho \omega_0} \right) \left(N_0 + \frac{1}{2} \mp \frac{1}{2} \right) \frac{D_{3D}(E \pm \hbar \omega_0)}{2} & N_0 &= \frac{1}{e^{\hbar \omega_0 / k_B T} - 1} \quad (\text{ODP}) \end{aligned}$$